

PROPERTIES AND CONSEQUENCES OF THORN-INDEPENDENCE

ALF ANGEL ONSHUUS

ABSTRACT. In the last couple of decades, independence relations have become one of the central parts of model theory. Areas such as classification theory, stability theory and simplicity theory use the notion of “forking independence” defined by Shelah as a main tool for their development. However, as powerful and useful as forking-independence is, it lacks generality. O-minimal structures give an example of a whole class of structures for which forking independence does not work. This, we believe, is the main cause why even though the results in o-minimal theory seem very similar to those in the theory of strongly minimal sets (strongly minimal sets are in some ways the smallest structures for which forking defines an independence relation) the proofs of analogous results have been very different in both areas.

We begin by developing a new notion of independence ($\mathfrak{p}$ -independence, read “thorn”-independence) that arises from a family of ranks suggested by Scanlon ($\mathfrak{p}$ -ranks). We prove that in a large class of theories (it includes simple theories and o-minimal theories) this notion has many of the properties needed for an adequate geometric structure.

Finally, we analyze the behavior of $\mathfrak{p}$ -forking in some theories where an independence relation had already been studied by other authors. We prove that $\mathfrak{p}$ -independence agrees with the usual independence notions in stable, supersimple and o-minimal theories. Furthermore, we give some evidence that the equivalence between forking and $\mathfrak{p}$ -forking in simple theories might be closely related to one of the main open conjectures in simplicity theory, the stable forking conjecture. In particular, we prove that in any simple theory where the stable forking conjecture holds, $\mathfrak{p}$ -independence and forking independence agree.

1. INTRODUCTION

1.1. Overview. In the last decades independence notions have become one of the central ideas in model theory. Especially in the last three decades independence notions and ranks have been the key factor first in Shelah’s classification theory and later in understanding the geometry in the models of theories such as stable, o-minimal and in the last decade simple theories. However, even when many of the results on the topology of o-minimal theories were similar to some of those in geometric stability theory, they were developed almost independently from each other mainly because of the difference between the definitions of the respective independence notions.

In sections 2 through 4 we develop a new notion of independence ($\mathfrak{p}$ -independence –read “thorn” independence–) and a rank that is associated to this independence notion ($\mathfrak{p}$ -rank). We prove that in a large class of theories which we shall call “rosy” this notion defines a geometric independence relation. This class of theories includes simple and o-minimal theories. It also includes theories for which there was no previously known independence relation; for example models of the theory defined by Casanovas and Wagner in [CW02]¹ are rosy.

In section 5 we study the relation between the classical independence notions which we had in simple and o-minimal theories and $\mathfrak{p}$ -independence. We prove that $\mathfrak{p}$ -independence agrees with the classical independence notion in o-minimal, stable and all known cases of simple theories. This provides a unified approach in two areas where, until now, the methods and proofs have been different (even though, as mentioned above, they did go in the same general direction and similar results were obtained). It also gives an alternative (and, in our point of view simplified) definition of forking in stable theories which might give further insight in stability theory.

1.2. Notation and Conventions. We assume the reader is familiar with the terminology and the basic results of model theory and, more specifically, stability and simplicity theory.

As it is common in stability theory, given a complete theory T we will fix a universe $\mathcal{C}$ called a “monster” model of T : we choose some saturated model $\mathcal{C}$ of cardinality κ and assume all sets, types and models we talk about have cardinality less than κ and live inside $\mathcal{C}$. In particular, by *models of T* we mean an elementary submodel of $\mathcal{C}$. Any automorphism will be understood to be a $\mathcal{C}$ -automorphism. Following [Hod93], an *equivalence formula* of $\mathcal{C}$ is a formula $\phi(x, y)$ in the language of T that defines an equivalence relation in $\mathcal{C}$. Unless otherwise specified, we will work inside $\mathcal{C}^{eq}$ in the sense of [She90]. By convention lower case letters a, b, c, d will in general represent tuples (of imaginaries) and upper case letters will represent sets. Greek letters such as $\delta, \sigma, \psi, \phi$ will be used for formulas.

2. $\mathfrak{p}$ -FORKING

2.1. Definitions. We will start by defining the notions that we will work with throughout this paper.

Definition 2.1. A formula $\delta(x, a)$ strongly divides over A if $tp(a/A)$ is non-algebraic and $\{\delta(x, a')\}_{a' \models tp(a/A)}$ is k -inconsistent for some $k \in \mathbb{N}$.

We will say that $\delta(x, a)$ $\mathfrak{p}$ -divides over A if we can find some tuple c such that $\delta(x, a)$ strongly divides over Ac .

Finally, a formula $\mathfrak{p}$ -forks over A if it implies a (finite) disjunction of formulae which $\mathfrak{p}$ -divide over A .

¹In this paper, Casanovas and Wagner construct a theory which is not simple and provides the first example of a theory without the strict order property which does not eliminate hyperimaginaries

Remark 2.1.1. Suppose a formula $\delta(x, a)$ strongly divides over some set C ; let $p(x, C) := tp(a/C)$. By definition there is some $k \in \mathbb{N}$ such that $\{\delta(x, a')\}_{a' \models tp(a/C)}$ is k -inconsistent.

Another way of saying this is that for all $x_1, x_2, \dots, x_k$

$$p(x_1, C) \cup p(x_2, C) \cup \dots \cup p(x_k, C) \models \neg \left(\bigwedge_{i=1}^k \phi(x, a_i) \right).$$

By compactness, there is some formula $\theta(y, c) \in tp(a/C)$ such that $\{\delta(x, a')\}_{a' \models \theta(y, c)}$ is k -inconsistent.

We will say the type $p(x)$ $\mathfrak{p}$ -divides over A if there is a formula in $p(x)$ which $\mathfrak{p}$ -divides over A ; similarly for $\mathfrak{p}$ -forking. We say that a is $\mathfrak{p}$ -independent of b over A , denoted $a \downarrow_A^{\mathfrak{p}} b$, if $tp(a/Ab)$ does not $\mathfrak{p}$ -fork over A .

Note that even though $\mathfrak{p}$ -dividing and dividing have similar definitions, the fact that we ask for k -inconsistency for a set of formulas for which the parameters vary in a definable class as opposed to an indiscernible sequence is a significant modification. In particular, as we will prove later, many theories with the strict order property will behave nicely under this new definition.

2.1.1. First Results.

Lemma 2.1.2. Let A, B, C be subsets (of $\mathcal{C}$) such that $A \subseteq B \subseteq C$ and let a, b be tuples. Then

- (1) Extension: Given a type p over B which does not $\mathfrak{p}$ -fork over A , we can extend p to a type q over C which does not $\mathfrak{p}$ -fork over A .
- (2) Monotonicity: If a formula $\delta(x, b)$ $\mathfrak{p}$ -forks over B , then it $\mathfrak{p}$ -forks over A .
- (3) Partial right transitivity: If $tp(a/C)$ does not $\mathfrak{p}$ -fork (divide) over A then $tp(a/C)$ does not $\mathfrak{p}$ -fork (divide) over B and $tp(a/B)$ does not $\mathfrak{p}$ -fork (divide) over A .
- (4) If $b \downarrow_A^{\mathfrak{p}} a$ and $tp(a/A)$ is non-algebraic, then $tp(a/Ab)$ is not algebraic.
- (5) If $b \downarrow_A^{\mathfrak{p}} a$ and $tp(b/A)$ is non-algebraic, then $tp(b/Aa)$ is not algebraic.
- (6) If $p(x, b)$ is a type over Ab which does not $\mathfrak{p}$ -fork over A and $\theta(x, y)$ is a formula such that $p(x, b) \models \exists y \theta(x, y)$. Then $p(x, b) \cup \{\theta(x, y)\}$ does not strongly divide over A .
- (7) Base Extension: If $a \downarrow_A^{\mathfrak{p}} c$, then for any tuple d there is some $d' \models tp(d/Ac)$ such that $a \downarrow_{Ad'}^{\mathfrak{p}} c$.

Proof.

1) Let $\Gamma(x) := \{\psi(x) \mid \psi(x) \in \mathcal{L}(x) \text{ and } \neg\psi(x) \text{ } \mathfrak{p}\text{-divides over } A\}$. As in the proof of extension for forking, we first need to prove that the type $p(x) \cup \Gamma(x)$ is consistent. Suppose this was not the case. By compactness there would be a finite subset $\psi_1(x), \dots, \psi_n(x)$ in $\Gamma(x)$ such that $p(x) \models \bigvee_{i=1}^n \neg\psi_i(x)$.

By definition $p(x)$ would $\mathfrak{p}$ -fork over A . Let q' be a complete extension of $p(x) \cup \Gamma(x)$ to some set containing C , and let q be the restriction of q' to C . By definition q does not $\mathfrak{p}$ -fork over A .

2) In the definition of $\mathfrak{p}$ -dividing we are allowed to add parameters to get strong division, so if any formula $\mathfrak{p}$ -divides over an extension of A then it $\mathfrak{p}$ -divides over A . The result for $\mathfrak{p}$ -forking follows.

3) The first implication follows from monotonicity and the other follows from the fact that we are considering fewer formulas.

4) Let $p(x, b) = tp(a/Ab)$. Assuming $p(x, b)$ does not strongly divide and using non-algebraicity of $tp(b/A)$, we can find distinct realizations $b_1, b_2, \dots, b_n, \dots$ of $tp(b/A)$ such that $\bigcup_i p(x, b_i)$ is consistent, realized by some a' . However, as $tp(b_1/A) = tp(b/A)$ we can assume $a'b_1 = ab$. However, $p(x, b)$ is a complete type, so $b_i \models tp(b/Aa)$ showing that $tp(b/Aa)$ is non algebraic.

5) We will prove the contrapositive of the statement. Let $tp(b/Aa)$ be algebraic and let $b = b_1, b_2, \dots, b_n$ be all the elements that satisfy $tp(b/Aa)$. Then

$$tp(b/Aa) \models (x = b_1) \vee (x = b_2) \vee \dots \vee (x = b_n).$$

If $tp(b/A)$ is non-algebraic, then $x = b_i$ strongly divides (and therefore $\mathfrak{p}$ -divides) over A for all $i \leq n$; by definition $tp(b/Aa)$ $\mathfrak{p}$ -forks over A and $b \not\downarrow_A^{\mathfrak{p}} a$.

6) Let us suppose there is some $\delta(x, b) \in p(x, b)$ such that $\delta(x, b) \wedge \theta(x, y)$ strongly divides over A and suppose $\delta(x, b) \wedge \exists y \theta(x, y)$ does not strongly divide over A . By definition (and compactness) there are some distinct $b_1, \dots, b_n, \dots$ such that $b_i \models tp(b/A)$ and $\bigwedge_{i \in \mathbb{N}} \delta(x, b_i) \wedge \exists y \theta(x, y)$ is consistent, satisfied by some element a . Let $c \models \theta(a, y)$. Then $(a, c) \models \delta(x, b_i) \wedge \theta(x, y)$ for all i , contradicting our assumptions.

7) By extension we can find some $a' \models tp(a/Ac)$ such that $a' \downarrow_A^{\mathfrak{p}} cd$. Let d' be the image of an automorphism which sends a' to a fixing Ac , so that $a \downarrow_A^{\mathfrak{p}} cd'$. By partial right transitivity, $a \downarrow_{Ad'}^{\mathfrak{p}} c$. $\square$

Remark 2.1.3. *Condition 5 above does not hold for $\mathfrak{p}$ -dividing in place of $\mathfrak{p}$ -forking, even in the theory in the language of equality that states that there are infinitely many elements. In fact, if A is the unordered pair $\{a, b\}$ where $a \leq b$, then $tp(a/A)$ is algebraic, but $x \in A$ does not $\mathfrak{p}$ -divide over the empty set.*

Proof. We will prove the remark by contradiction. Let B be any set such that $tp(A/B)$ is non-algebraic and $x \in A$ strongly divides over B . If $tp(a/Bb)$ was non-algebraic, then $\{x \in A'\}_{A' \models tp(A/B), b \in A'}$ would be an infinite consistent set of formulas which would contradict strong division. This means that $b \in \text{acl}(Ba)$; by symmetry, $a \in \text{acl}(Bb)$. In this particular theory we may assume that B is a subset of M (as opposed to M^{eq})². But $tp(A/B)$ is non-algebraic, so $tp(b/B)$ and $tp(a/B)$ must be both be non-algebraic. By

²This is because the theory of equality has “elimination of imaginaries”, a property we will talk about further on

Steinitz exchange principle (see for example [Pil96], definition 2.1.1), there must be some element in B algebraic over a . However, $\text{acl}(a) = \{a\}$, so $a \in B$ which contradicts a being non-algebraic over B . $\square$

This shows that, unlike the case of simple theories, even when we are only considering the language of equality there is a difference between p -dividing and p -forking.

The following results go in the same direction as the ones above (showing properties that p -forking has in a general theory), but the proofs are a little more elaborated.

Theorem 2.1.4. *Given a tuple a and sets $A \subset B$, then $tp(a/B)$ p -forks over A if and only if $tp(a/\text{acl}(B))$ p -forks over A . In other words, $a \downarrow_A^b B$ if and only if $a \downarrow_A^b \text{acl}(B)$.*

Proof. Monotonicity implies that whenever $tp(a/B)$ p -forks over A , $tp(a/\text{acl}(B))$ p -forks over A . For the other direction, suppose $\psi(x, c) \in tp(a/\text{acl}(B))$ is such that $\psi(x, c)$ p -forks over A . Let $\phi_i(x, d_i)$ be such that

$$\psi(x, c) \Rightarrow \phi_i(x, d_1) \vee \phi(x, d_2) \vee \cdots \vee \phi(x, d_m)$$

and for any i , $\phi_i(x, d_i)$ p -divides over A . Let $c_1, \dots, c_n$ be the conjugates of c over B and for $0 \leq j \leq n$ let σ_j be some B -automorphism such that $\sigma_j(c) = c_j$. This means that $tp(a/B) \models \bigvee_{j=1}^n \psi(x, c_j)$ and for any such j $\psi_j(x, c_j) \Rightarrow \bigvee_{i=1}^m \phi_i(x, \sigma_j(d_i))$. Combining this two, we get

$$tp(a/B) \models \bigvee_{j=1}^n \bigvee_{i=1}^m \phi_i(x, \sigma_j(d_i))$$

and by definition $tp(a/B)$ p -forks over A . $\square$

We can also prove without any assumptions on the theory a version of partial left transitivity:

Lemma 2.1.5. *Partial Left Transitivity: If $a \downarrow_A^b c$ and $b \downarrow_{Aa}^b c$ then $ab \downarrow_A^b c$.*

Proof. We begin with a claim.

Claim 2.1.6. *It is enough to show that $tp(ab/Ac)$ does not p -divide over A .*

Proof. Suppose $a \downarrow_A^b c$, $b \downarrow_{Aa}^b c$ and $ab \not\downarrow_A^b c$. Choose $\phi(x, y, c) \in tp(ab/Ac)$ which implies $\bigvee_{i < n} \psi_i(x, y, c_i)$, where each ψ_i p -divides over A . Put $\bar{c} := (c_1 : i < n)$. By extension we can find some $a' \models tp(a/Ac)$ such that $a' \downarrow_A^b c\bar{c}$. Let b_2 be the image of b under an automorphism fixing Ac which sends a to a' , so that $tp(ab/Ac) = tp(a'b_2/Ac)$ and $b_2 \downarrow_{Aa'}^b c$. By extension again we can also find some $b' \models tp(b_2/Aa'c)$ such that $b' \downarrow_{Aa'}^b c\bar{c}$. So we have tuple $a'b'$ satisfying the hypothesis of the theorem. However, $\phi(x, y, c) \in tp(a'b'/Ac)$ and $\phi(x, y, c) \Rightarrow \bigvee \psi_i(x, y, c_i)$, so $\psi_i(x, y, c_i) \in tp(a'b'/Ac\bar{c})$ for some $i < n$, and $tp(a'b'/Ac\bar{c})$ p -divides over A . $\square$

Let us suppose then that $tp(ab/Ac) \models \phi(x, y, c)$ which $\mathfrak{p}$ -divides over A strongly dividing, say, over Ad . By base extension, we can find some $d' \models tp(d/Ac)$ such that $a \downarrow_{Ad'}^{\mathfrak{p}} c$, so we will assume $a \downarrow_{Ad}^{\mathfrak{p}} c$. By definition of strong dividing we know that $\{\phi(x, y, z)\}_{z \models tp(c/Ad)}$ is k -inconsistent and $tp(c/Ad)$ is non algebraic; by 2.1.2(4) so is $tp(c/Ada)$. This means that

$$\{\phi(a, y, z)\}_{z \models tp(c/Ada)}$$

is still k -inconsistent and $tp(c/Ada)$ is non-algebraic. By definition $tp(b/Aac)$ strongly divides over Aad which means it $\mathfrak{p}$ -divides over Aa , a contradiction. $\square$

As in simple theories, this new notion of bifurcation does have some relation with independent sequences which will be stated in the following lemma. However, unlike in simple theories this relation does not seem to be fundamental for the development of the theory, perhaps because the main function of indiscernible sequences in simple theories is to provide some uniformity when witnessing division; such uniformity is provided by the definability of the parameters in the the definition of $\mathfrak{p}$ -dividing.

Lemma 2.1.7. *Let a, b be elements and A a set. Let $p(x, b) = tp(a/Ab)$. Then the following conditions are equivalent:*

- (1) $tp(a/Ab)$ does not $\mathfrak{p}$ -divide over A
- (2) For any $B \supseteq A$ such that b is not algebraic over B , there is some tuple $a' \models tp(a/Ab)$ and some infinite Ba' -indiscernible sequence I containing b .

Proof. ($\Leftarrow$) We will proceed by contradiction. Assume $tp(a/Ab)$ $\mathfrak{p}$ -divides over A . By definition we have a B with $b \notin acl(B)$ and a δ , such that $\models \delta(a, b)$ and $\{\delta(x, b')\}_{b' \models tp(b/B)}$ is k -inconsistent. Let I be any B -indiscernible sequence. Since the underlying set of I is a subset of $\{b' \mid b' \models tp(b/B)\}$, we have that $\{\delta(x, b')\}_{b' \in I}$ is k -inconsistent. Let $a' \models tp(a/Ab)$, $a' \models \delta(x, b)$. By k -inconsistency we have $\not\models \delta(a', b')$ for all but finite $b' \in I$ which implies that I is not Ba' indiscernible.

($\Rightarrow$) Let A, b, a be such that $tp(a/Ab)$ does not $\mathfrak{p}$ -divide over A . If b is algebraic over A , condition 2 in the lemma holds immediately so there is nothing to prove. We will assume then that b is not algebraic over A .

Let B be any set containing A with $b \notin acl(B)$ and let $q(y) = tp(b/B)$. We know that for any $\delta(x, b) \in tp(a/Ab)$ the set $\{\delta(x, b')\}_{b' \models tp(b/B)}$ is not k -inconsistent for any k . By compactness

$$\bigcup_{i \in \omega} q(y_i) \cup \{p(x, y_i) : i \in \omega\}$$

is consistent, and realized by a', I , say. For any finite set Δ of formulas we can find by Ramsey's theorem some infinite $I_\Delta \subseteq I$ such that I_Δ is Δ -indiscernible over Ba' . By compactness we can find a Ba' -indiscernible sequence J such that $a'b' \models p(x, y) \cup q(y)$ for any $b' \in J$. If σ is a B -automorphism with $\sigma(b') = b$ then $\sigma(a'), \sigma(J)$ will do. $\square$

2.2. Existence. Given some independence relation, we say that such a relation satisfies *existence* if for any tuple a and any set A , a is independent with A over A . This is a very useful property for simple theories. Until this point we have studied the behavior of $\mathfrak{p}$ -forking in the most general context. In this section we will only consider theories for which $\mathfrak{p}$ -forking satisfies existence: given a tuple a and a set A , $tp(a/A)$ does not $\mathfrak{p}$ -fork over A .

As we mentioned before, unlike in simple theories there is a difference between $\mathfrak{p}$ -forking and $\mathfrak{p}$ -dividing. As we shall see in most of the proofs, we can usually work around this problem since extension provides a way to reduce most of the proofs down to $\mathfrak{p}$ -dividing. It would be nice however to have some idea of the relation between the parameters that we need for $\mathfrak{p}$ -forking and those that are used in the corresponding $\mathfrak{p}$ -dividing formulas, so we can tell how much do we have to extend a $\mathfrak{p}$ -forking type before achieving $\mathfrak{p}$ -division. The next lemma gives a partial answer to this question.

Lemma 2.2.1. *If $\delta(x, a)$ is consistent and $\mathfrak{p}$ -forks over A , as witnessed by a disjunction $\bigvee_{i=1}^n \psi(x, a_i)$ implied by $\delta(x, a)$, such that $\psi_i(x, a_i)$ strongly divides over Aa_i , then a_i is algebraic over $Aa\bar{c}$ where $\bar{c} = (c_i : i < n)$. Even more, for at least one i , a_i is algebraic over Aac_i . (We are assuming that there are no “extra” ψ_i ’s: i.e. that $\delta(x, a) \not\models \bigvee_{j \in I} \psi(x, a_j)$ for any $I \subsetneq \{1, 2, \dots, n\}$.)*

Proof. Let b be any element such that $b \models \delta(x, a)$. By existence and definition of $\mathfrak{p}$ -forking we can extend $tp(b/Aa)$ to $tp(b/Aa\bar{a}_i\bar{c})$ so that $b \perp_{Aa}^{\mathfrak{p}} \bar{a}_i\bar{c}$. In particular, by partial transitivity $b \perp_{Aac_i}^{\mathfrak{p}} a_i$.

We know that $b \models \psi_i(x, a_i)$ for some i ; for any $a'_i \models tp(a_i/Abac_i)$ we have $\mathcal{C} \models \psi(b, a'_i)$ and $a'_i \models tp(a_i/Ac_i)$. By the definition of strong dividing there cannot be infinitely many such a'_i ’s (b would witness the consistency). Thus $tp(a_i/Abc_i)$ must be algebraic and therefore so is $tp(a_i/Abac_i)$. But we know that $b \perp_{Aac_i}^{\mathfrak{p}} a_i$, so by lemma 2.1.2(4) this means that $tp(a_i/Aac_i)$ is algebraic.

To finish the proof, we just have to be careful when extending $tp(b/Aa)$ to Aaa_i ; let $\delta(x, a, a_i)$ be $\delta(x, a) \wedge \neg \psi_i(x, a_i)$ so $\delta(x, a, a_i)$ implies the disjunction $\bigvee_{j \neq i} \psi_j(x, a_j)$. Since k -inconsistency is preserved, either there is some a_j algebraic over Aa_i or $\delta(x, a, a_i)$ $\mathfrak{p}$ -forks over Aa_i in which case we can repeat the process and get some a_j algebraic over $Aa_i c_j$. Either way we get a_j algebraic over $Aa_i c_j$ for some j . However, a_i is algebraic over Aa_i so a_j is algebraic over $Aa_i c_j$. $\square$

Theorem 2.2.2. *Let $p(x, b)$ be a type over Ab which is non- $\mathfrak{p}$ -forking over A and let $\theta(x, y)$ be a formula such that $p(x, b) \models \exists y \theta(x, y)$. Then $p(x, b) \cup \{\theta(x, y)\}$ does not $\mathfrak{p}$ -fork over A .*

Proof. Suppose that it does $\mathfrak{p}$ -fork over A so that $p(x, b) \cup \{\theta(x, y)\} \models \bigvee_{i=1}^n \psi_i(x, y, a_i)$ where $\psi_i(x, y, a_i)$ $\mathfrak{p}$ -divides over A by strongly dividing over Aa_i .

By hypothesis $p(x, b)$ does not $\mathfrak{p}$ -fork over A , so using extension we can choose some $a \models p(x, b)$ such that $a \downarrow_A^{\mathfrak{p}} b\bar{a}_i\bar{c}_i$ which implies by partial transitivity that $a \downarrow_{Ac_i}^{\mathfrak{p}} a_i$. On the other hand, using existence and extension we can choose some $c \models \theta(a, y)$ such that $c \downarrow_{Aa}^{\mathfrak{p}} b\bar{a}_i\bar{c}_i$. Using partial transitivity again we get $c \downarrow_{Aac_i}^{\mathfrak{p}} a_i$.

We know that $(a, c) \models p(x, b) \cup \theta\{(x, y)\}$ so $(a, c) \models \psi_j(x, y, a_j)$ for some j . By definition, a_j is algebraic over $Aacc_j$. However, using lemma 2.1.2 with $c \downarrow_{Aac_j}^{\mathfrak{p}} a_j$ we get that a_j is algebraic over Aac_j . Using 2.1.2 one more time will give us a_j algebraic over Ac_j , contradicting the definition of strong dividing. $\square$

Definition 2.2. A notion of independence $\downarrow^0$ has the strong extension property if whenever $a \downarrow_A^0 b$ then for any c there is $c' \models tp(c/Aa)$ with $ac' \downarrow_A^0 b$, or equivalently, for any c there is $b' \models tp(b/Aa)$ with $ac \downarrow_A^0 b'$.

Corollary 2.2.3. $\mathfrak{p}$ -independence has the strong extension property.

Proof. It is enough to prove that if $p(x, b)$ is a non forking extension of $p \upharpoonright A$ and $q(x, y)$ is a consistent type over A containing $p \upharpoonright A$ then $p(x, b) \cup q(x, y)$ does not $\mathfrak{p}$ -fork over A . This follows from theorem 2.2.2, as consistency of $(p \upharpoonright A) \cup q$ implies consistency of $p \cup q$. $\square$

Definition 2.3. A sequence of elements is a $\mathfrak{p}$ -Morley sequence over A if it is indiscernible and $\mathfrak{p}$ -independent over A .

Claim 2.2.4. Let $p(x)$ be a complete type over $B \supset A$ which does not $\mathfrak{p}$ -fork over A . Then there is a $\mathfrak{p}$ -Morley sequence over A with all of its elements realizing $p(x)$.

The claim is true for any independent notion that has extension. The proof is exactly the same as the one given in the simple theoretic context (see [Wag00]). We will prove later that in the theories that will actually interest us, $\mathfrak{p}$ -Morley sequences will provide an alternative definition of $\mathfrak{p}$ -forking.

3. $\mathfrak{p}$ -RANK AND ROSY THEORIES

3.1. Definition of $\mathfrak{p}$ -rank. We will now define a notion of rank that will code $\mathfrak{p}$ -forking.

Definition 3.1. Given a formula ϕ , a set Δ of formulas in the variables $x; y$, a set of formulae Π in the variables $y; z$ (with z possibly of infinite length) and a number k , we define $p(\phi, \Delta, \Pi, k)$ inductively as follows:

- (1) $p(\phi, \Delta, \Pi, k) \geq 0$ if ϕ is consistent.
- (2) For λ limit ordinal, $p(\phi, \Delta, \Pi, k) \geq \lambda$ if and only if $p(\phi, \Delta, \Pi, k) \geq \alpha$ for all $\alpha < \lambda$

- (3) $p(\phi, \Delta, \Pi, k) \geq \alpha + 1$ if and only if there is a $\delta \in \Delta$, some $\pi(y; z) \in \Pi$ and parameters c such that
- (a) $p(\phi \wedge \delta(x, a), \Delta, \Pi, k) \geq \alpha$ for infinitely many $a \models \pi(y; c)$
 - (b) $\{\delta(x, a)\}_{a \models \pi(y; c)}$ is k -inconsistent

As usual, for a type p , we define

$$p(p, \Delta, \Pi, k) = \min \{ p(\phi(x), \Delta, \Pi, k) \mid \phi(x) \in p \}.$$

Remark 3.1.1. We can give a definition directly for types, changing all instances of ϕ for some type p which is the case we will usually use. However, when dealing with types, we will use an alternative version of condition 3(a). Let p be a complete type over A and let us assume that we are witnessing the rank going up as in condition 3, with c being a tuple containing all the parameters in π . We can always find some non-algebraic complete type $q(y)$ over Ac containing π such that for any $a' \models q(y)$, $p(\phi \wedge \delta(x, a'), \Delta, \Pi, k) \geq \alpha$. Therefore, we can change condition 3(a) by

3(a)'. $p(p \cup \{\delta(x, a)\}, \Delta, \Pi, k) \geq \alpha$ where $a \models \pi$ and $tp(a/Ac)$ is non-algebraic.

Remark 3.1.2. Given any type $p(x; y)$ (not necessarily complete), formulas ϕ, π and integers k, n , the set

$$\{b \mid p(p(x, b), \phi, \pi, k) \geq n\}$$

is type definable.

This has nice consequences for the structure of the p -rank. For example, compactness implies that $p(p \cup \{\delta(x, a)\}, \Delta, \Pi, k)$ is finite whenever it is defined.

Proof. Notice that

$$p(p(x, b), \phi, \pi, k) \geq n$$

is witnessed by a tree where each of the nodes is a formula $\phi(x, a)$, its height is n , for each $i \leq n$ there is some c_i such that the level i contains all $\phi(x, a)_{a \models \pi(x, c_i)}$, the union any k of such formulas (all in the same level) is inconsistent and the union of any branch is consistent with p . All these properties can be described by formulas, and we get type definability. $\square$

3.2. Properties of the p -rank.

Theorem 3.2.1. This thorn rank has the following properties:

- (1) Monotonicity: If $\Delta \subseteq \Delta'$, $p \supseteq p'$, and $\Pi \subseteq \Pi'$ then

$$p(p', \Delta', \Pi', k) \geq p(p, \Delta, \Pi, k).$$

- (2) Transitivity: If $p \subseteq q \subseteq r$, then $p(r, \Delta, \Pi, k) = p(p, \Delta, \Pi, k)$ if and only if $p(r, \Delta, \Pi, k) = p(q, \Delta, \Pi, k)$ and $p(q, \Delta, \Pi, k) = p(p, \Delta, \Pi, k)$.

- (3) Additivity: $p((\theta \vee \psi), \Delta, \Pi, k) = \max \{ p(\theta, \Delta, \Pi, k), p(\psi, \Delta, \Pi, k) \}$

Proof. .

1) We will prove by induction on λ that if $\lambda \leq \text{p}(p, \Delta, \Pi, k)$ then $\text{p}(p', \Delta', \Pi', k) \geq \lambda$. For $\lambda = 0$, the proof is clear, as is the induction step for the case when λ is a limit ordinal. Now, suppose it is true for $\lambda = \alpha$ and let us assume that $\text{p}(p, \Delta, \Pi, k) \geq \alpha + 1$. Let ψ be any formula in p' ; by hypothesis $\psi \in p$ so $\text{p}(\psi, \Delta, \Pi, k) \geq \alpha + 1$. We can therefore find a $\phi \in \Delta$ and a formula π in Π such that $\text{p}(\psi \wedge \phi(x, a), \Delta, \Pi, k) \geq \alpha$ for infinitely many $a \models \pi$ and $\{\phi(x, a)\}_{a \models \pi}$ is k -inconsistent. By induction hypothesis $\text{p}(\psi \wedge \phi(x, a), \Delta', \Pi', k) \geq \alpha$. But $\psi \in p'$, $\phi \in \Delta'$ and $\pi \in \Pi'$ so ϕ actually witnesses $\text{p}(\psi, \Delta', \Pi', k) \geq \alpha + 1$ for all $\psi \in p'$.

2) By monotonicity, $\text{p}(r, \Delta, \Pi, k) \leq \text{p}(q, \Delta, \Pi, k) \leq \text{p}(r, \Delta, \Pi, k) \geq$; transitivity follows from transitivity for equality.

3) By monotonicity we have

$$\text{p}((\theta \vee \psi), \Delta, \Pi, k) \geq \max\{\text{p}(\theta, \Delta, \Pi, k), \text{p}(\psi, \Delta, \Pi, k)\}.$$

For the other direction we will prove by induction on α that if

$$\text{p}((\theta \vee \psi), \Delta, \Pi, k) \geq \alpha$$

then either $\text{p}(\theta, \Delta, \Pi, k) \geq \alpha$ or $\text{p}(\psi, \Delta, \Pi, k) \geq \alpha$. Once again, the only difficult step is the induction step. Let us assume it is true for α and that $\text{p}((\theta \vee \psi), \Delta, \Pi, k) \geq \alpha + 1$. We can then find a $\delta \in \Delta$ and a formula $\pi \in \Pi$ such that

$$\text{p}((\theta \vee \psi) \wedge \delta(x, a_i), \Delta, \Pi, k) \geq \alpha$$

for infinitely many $a_i \models \pi$ and $\{\phi(x, a_i)\}_{a_i \models \pi}$ is k -inconsistent. By induction, for any such a_i either $\text{p}(\theta \wedge \delta(x, a_i), \Delta, \Pi, k) \geq \alpha$ or $\text{p}(\psi \wedge \delta(x, a_i), \Delta, \Pi, k) \geq \alpha$; but then for one of ψ or θ (let us assume θ) we have infinitely many a_i 's such that $\text{p}(\theta \wedge \delta(x, a_i), \Delta, \Pi, k) \geq \alpha$ and by definition $\text{p}(\theta, \Delta, \Pi, k) \geq \alpha + 1$. $\square$

Corollary 3.2.2. Extension for fixed Δ and Π :

For any partial type ξ defined over a set A , finite sets of formulas Δ and Π and any k , we can extend ξ to a complete type p over A such that $\text{p}(p, \Delta, \Pi, k) = \text{p}(\xi, \Delta, \Pi, k)$.

Proof. It is just the usual application of additivity, the definition of p -rank for types and Zorn's lemma as it is used in simple theories (see [Kim96]). $\square$

One of the properties that we will prove here is that p -independence and p -ranks are related in the same way that forking and the D -ranks are. The next theorem is one of the directions of the relation we will prove.

Theorem 3.2.3. *Let p be a type over $B \supseteq A$ such that for every ϕ, Π, k ,*

$$\text{p}(p \upharpoonright A, \phi, \Pi, k) = \text{p}(p, \phi, \Pi, k)^3.$$

Then p does not p -fork over A .

³By $\text{p}(p, \phi, \Pi, k)$ we really mean $\text{p}(p, \{\phi\}, \Pi, k)$.

Proof. Suppose p $\mathfrak{b}$ -forks over A . So

$$p \models \bigvee_{i < n} \phi_i(x, b_i)$$

where each ϕ_i $\mathfrak{b}$ -divides over A . By definition there is some $\theta_i(y, z)$, an element d_i and some finite k_i such that $\{\phi(x, b'_i)\}_{b'_i \models \theta_i(y, d_i)}$ is k_i -inconsistent and $tp(b_i/Ad_i)$ is a non-algebraic type containing $\theta(y, d_i)$. We can always take the maximum of the k_i 's and assume they are all equal.

Now, as in the proof for simple theories (see [Kim96]), we construct a formula

$$\psi(x, y_1, \dots, y_n, z) := \bigvee_{i < n} (\phi_i(x, y_i) \wedge y_i = z)$$

which uniformizes the ϕ_i in the following sense: if $\bar{y} = (y_1, \dots, y_n, w)$ then for any $i \leq n$, $w = y_i$ implies $\psi(x, \bar{y}) \iff \phi_i(x, y_i)$. On the other hand, if we define a new formula $\theta'_i(\bar{y}, z) := \{w = y_i\} \cup \theta_i(y_i, z)$ we have that for any tuple $\bar{b} = \langle b_1, b_2, \dots, b_n, l \rangle$, and for any d , $\bar{b} \models \theta'_i(y, d)$ if and only if $b_i \models \theta_i(y, d)$ and $l = b_i$. Thus, if we define $c_i := \langle b_1, \dots, b_n, b_i \rangle$ we get $p \models \bigvee_{i < n} \psi(x, c_i)$ and $\{\psi(x, c_i)\}_{c_i \models \theta'_i(y, d)}$ is k -inconsistent. By additivity we can add at least one of the ϕ_i 's without changing the rank, so we can extend p to some type q over $Bc_1 \dots c_n$ such that $\mathfrak{p}(p, \psi, \{\theta'_1, \dots, \theta'_n\}, k) = \mathfrak{p}(q, \psi, \{\theta'_1, \dots, \theta'_n\}, k)$ and q implies one of the $\phi_i(x, b_i)$ and thus one of the $\psi(x, c_i)$. Whichever one it is (we can assume it is the first one without loss of generality), we know that $\{\psi(x, t)\}_{t \models \theta'_1(y, b)}$ is k -inconsistent and $tp(\bar{b}/Ad)$ is a non-algebraic type containing $\theta'_1(y, d)$. By definition of $\mathfrak{p}$ -rank

$$\mathfrak{p}(p \upharpoonright A, \psi, \{\theta'_1, \dots, \theta'_n\}, k) \geq \mathfrak{p}(q, \psi, \{\theta'_1, \dots, \theta'_n\}, k) + 1$$

and therefore

$$\mathfrak{p}(p, \psi, \{\theta'_1, \dots, \theta'_n\}, k) < \mathfrak{p}(p \upharpoonright A, \psi, \{\theta'_1, \dots, \theta'_n\}, k),$$

a contradiction. $\square$

Corollary 3.2.4. *If for any finite Δ, Π all the $\mathfrak{p}$ -ranks are defined, then we have existence (and all the results of subsection 3).*

Proof. Let $p(x)$ be a type over A . Then $p \upharpoonright A = p$ and for any finite set of formulas Δ, Π and any finite k ,

$$\mathfrak{p}(p \upharpoonright A, \phi, \Pi, k) = \mathfrak{p}(p, \phi, \Pi, k).$$

The theorem implies that $p(x)$ does not $\mathfrak{b}$ -fork over A . $\square$

4. ROSY THEORIES

As the corollary above suggests, there is a lot to say about theories that have ordinal-valued $\mathfrak{p}$ -ranks even if we limit ourselves to finite formulas and types in the definition. Doing this has the extra advantage that, if we limit ourselves to finite Δ and Π , definability of $\mathfrak{p}$ -rank and compactness give us that a $\mathfrak{p}$ -rank is defined if and only if it is finite.

From now on, we will study the class of theories such that $\mathfrak{p}(p, \Delta, \Pi, k)$ is finite for any type $p(x)$ in the language of the theory, any finite sets of formulas Δ and Π and any finite number k . We will call any such theory “rosy”⁴. We will prove that in any rosy theory $\mathfrak{p}$ -forking has a lot of the geometric properties we want for an independence notion. All theories in this section are assumed to be rosy.

4.1. Geometry of $\mathfrak{p}$ -independence in Rosy theories.

Theorem 4.1.1. Symmetry: *For any two elements a, b and any set A , $a \downarrow_A^{\mathfrak{p}} b$ if and only if $b \downarrow_A^{\mathfrak{p}} a$.*

Proof. Let us suppose $a \downarrow_A^{\mathfrak{p}} b$ and let us assume that $\models \delta(b, a)$ where $\delta(x, a)$ $\mathfrak{p}$ -forks over A . Let $\delta(x, a) \Rightarrow \bigvee \psi_i(x, c_i)$ where each $\psi_i(x, c_i)$ strongly divides over Ad_i . By compactness we can prove the k -inconsistency using some formula π_i with parameters in Ad_i .

We define a sequence $\{a^j, \bar{c}^j, \bar{d}^j\}_{j \in \mathbb{N}}$ in the following way. Let $a^0 = a$, $\bar{c}^0 = \bar{c}_i$ and $\bar{d}^0 = \bar{d}_i$. Assuming we have defined the sequence up to $j = n$, let $a^{n+1} \models tp(a/Ab)$ be such that

$$a^{n+1} \downarrow_A^{\mathfrak{p}} ba^1 \bar{c}_i^1 \bar{d}_i^1 \dots a^n \bar{c}_i^n \bar{d}_i^n$$

(we know such a^{n+1} exists by extension) and let c_i^{n+1} and d_i^{n+1} be the images of c_i and d_i under an automorphism that sends a to a^{n+1} .

Once we have such a sequence, by right partial transitivity and monotonicity (2.1.2) we have that for all $n > m$,

$$a^n \downarrow_{Ad_i^m a^{m+1} \dots a^{n-1}}^{\mathfrak{p}} c_i^m.$$

Using induction and left partial transitivity (2.1.5) $n - m$ times we get

$$a^{m+1} a^{m+2} \dots a^n \downarrow_{Ad_i^m}^{\mathfrak{p}} c_i^m$$

so in particular it cannot strongly divide. But $tp(c_i^m / Ad_i^m)$ is non-algebraic so

$$tp(c_i^m / d_i^m a^{m+1} a^{m+2} \dots a^n)$$

is non-algebraic for any $n > m$; by local character of $\mathfrak{p}$ -forking and compactness

$$tp(c_i^m / d_i^m a^{m+1} a^{m+2} \dots)$$

would also be non-algebraic. By definition, $\psi_i(x, c_i^n)$ strongly divides over $Ad_i^n a^{n+1} \dots$ and $\psi_i(x, c_i^n)$ $\mathfrak{p}$ -divides over $Aa^{n+1} a^{n+2} \dots$.

⁴Rosy theories include simple and o-minimal theories but, as mentioned in the introduction, there are other rosy theories which do not fall into either of these categories.

Now, since $a^n \bar{c}_i^n \models tp(a\bar{c}_i/A)$, $a^n \models tp(a/Ab)$ we have that for all n

$$\delta(x, a^n) \Rightarrow \bigvee \psi_i(x, c_i^n)$$

and $b \models \delta(x, a^n)$. This implies $\delta(x, a^n) \in tp(b/Aa^n a^{n+1} a^{n+2} \dots)$ $\bar{\vdash}$ -forks over

$Aa^{n+1} a^{n+2} \dots$. By the proof of the previous theorem we know that this is witnessed by

$$\bar{\vdash}(tp(b/Aa^n a^{n+1} a^{n+2} \dots), \psi, \theta, k) < \bar{\vdash}(tp(b/Aa^{n+1} a^{n+2} \dots), \psi, \theta, k)$$

where θ is the one mentioned at the end of the proof of theorem 3.2.3 and ψ depends only on the ψ_i (so θ and ψ_i are the same for all n). This means that $\bar{\vdash}(tp(b/Aa^1 a^2 a^3 \dots), \psi, \theta, k)$ is infinite contradicting rosiness. $\square$

If we combine this result with those above, we get an analogue for most of the geometric properties of forking independence for simple theories (see [Wag00] 2.3.13). We will use the same names.

Corollary 4.1.2. *$\bar{\vdash}$ -independence defines an independence relation –in the sense described in [KP97]– in any rosy theory. More precisely, if we are working inside a theory T of finite $\bar{\vdash}$ -ranks, then the following properties hold for types in models of T .*

- (1) Existence: If $p \in S(A)$, then p does not $\bar{\vdash}$ -fork over A .
- (2) Extension: Every partial type over $B \supset A$ which does not $\bar{\vdash}$ -fork over A can be extended to a complete type $p(x)$ over B which does not $\bar{\vdash}$ -fork over A .
- (3) Reflexivity: $B \bar{\perp}_A^b B$ if and only if $B \subseteq acl(A)$.
- (4) Monotonicity: If p and q are types with $p \supseteq q$ and p does not $\bar{\vdash}$ -fork over A , then q does not $\bar{\vdash}$ -fork over A .
- (5) Finite Character: $C \bar{\perp}_A^b B$ if and only if $c \bar{\perp}_A^b B$ for any finite $c \subset C$.
- (6) Symmetry: $C \bar{\perp}_A^b B$ if and only if $B \bar{\perp}_A^b C$.
- (7) Transitivity: If $A \subset B \subset C$ then $b \bar{\perp}_A^b C$ if and only if $b \bar{\perp}_A^b B$ and $b \bar{\perp}_B^b C$.
- (8) If $a \bar{\perp}_A^b b$ and for some formula $\delta(x, y)$ $\delta(x, a)$ $\bar{\vdash}$ -forks over A , then $\delta(x, a)$ $\bar{\vdash}$ -forks over Ab .
- (9) Let $A \subset B$. If $a \bar{\perp}_A^b B$ then $a \bar{\perp}_A^b acl(B)$.

Proof. Properties 1,2,3,4,6,7,8 and 10 have already been proven or follow immediately as a corollary of symmetry. 5. is clear from the definitions.

To prove 9. we will first show that if $\delta(x, a)$ $\bar{\vdash}$ -divides over A , then it $\bar{\vdash}$ -divides over Ab . Take $B \supseteq A$ such that $\{\delta(x, a')\}_{a' \models tp(a/B)}$ is k -inconsistent and $tp(a/B)$ is non-algebraic. By base extension, we can find some $B' \models tp(B/Aa)$ such that $a \bar{\perp}_{B'}^b b$; note that $\delta(x, a)$ strongly divides over B' so $tp(a/B')$ is non-algebraic. By 2.1.2 $tp(a/B'b)$ is not algebraic and $\{\delta(x, a')\}_{a' \models tp(a/B'b)}$ is still k -inconsistent. By definition $\delta(x, a)$ $\bar{\vdash}$ -divides over Ab .

Suppose now that $\delta(x, a)$ $\mathfrak{p}$ -forks over A and $\delta(x, a) \Rightarrow \bigvee \phi_i(x, a_i)$ where each formula $\phi_i(x, a_i)$ $\mathfrak{p}$ -divides over A . By extension we can assume $a_i a \downarrow_A^{\mathfrak{p}} b$ which implies $a_i \downarrow_A^{\mathfrak{p}} b$. But we just proved that $\phi_i(x, a_i)$ $\mathfrak{p}$ -divides over Ab and thus $\delta(x, a)$ $\mathfrak{p}$ -forks over Ab . $\square$

As we mentioned before, in any rosy theory $\mathfrak{p}$ -forking is witnessed by the $\mathfrak{p}$ -ranks in the same way that forking was witnessed by the D -ranks. We already proved one of the implications in 3.2.3; the following theorem will prove the other one.

Theorem 4.1.3. *Let p be a type over $B \supseteq A$ such that p does not $\mathfrak{p}$ -fork over A . Then for all ϕ, θ, k $\mathfrak{p}(p \upharpoonright A, \phi, \theta, k) = \mathfrak{p}(p, \phi, \theta, k)$.*

Proof. Since whenever Δ and Π are finite $\mathfrak{p}(p, \Delta, \Pi, k)$ is finite for any type p and any k , it is enough to prove that for any natural number m , $\mathfrak{p}(p \upharpoonright A, \phi, \theta, k) \geq m$ implies $\mathfrak{p}(p, \phi, \theta, k) \geq m$. The case $m = 0$ is clear by the consistency of p . Suppose then that we have it for $m = n$ and that $\mathfrak{p}(p \upharpoonright A, \phi, \theta, k) \geq n + 1$. By definition we can find some b and some tuple c such that $\mathfrak{p}(p \upharpoonright A \cup \{\phi(x, b)\}, \phi, \theta, k) \geq n$, $b \models \theta(y, c)$, $\{\phi(x, b')\}_{b' \models \theta(y, c)}$ is k -inconsistent and $tp(b/Ac)$ is non-algebraic.

By 3.2.1 we can extend $p \upharpoonright A \cup \{\phi(x, b)\}$ to a type $q(x, b)$ over Ab such that

$$\mathfrak{p}(q(x, b), \phi, \theta, k) \geq n.$$

Now, if $a \models p$ and $a' \models q$, we have an automorphism that fixes A and sends a to a' , p to some type p' over B' which does not $\mathfrak{p}$ -fork over A and c to some c' which still witnesses the $\mathfrak{p}$ -division of $\phi(x, b)$. Since ϕ and θ have no parameters they are fixed under automorphisms so it is enough to show that $\mathfrak{p}(p', \phi, \theta, k) \geq n + 1$. We may therefore assume without loss of generality that $p = p'$, $a = a'$ and thus a satisfies both q and p . By extension we can extend $tp(b/Aa)$ to a type over Ba which does not $\mathfrak{p}$ -fork over Aa ; let b' be some realization of such extension. Let σ be some automorphism that sends b to b' , fixes fixing Aa and sends c to c' .

We may assume $b' = b$ and $b \downarrow_{Aa}^{\mathfrak{p}} B$. By hypothesis, $a \downarrow_A^{\mathfrak{p}} B$ and by transitivity, $tp(ba/B)$ does not $\mathfrak{p}$ -fork over A . This implies that $tp(a/Bb)$ does not $\mathfrak{p}$ -fork over Ab (by transitivity) and $tp(B/baA)$ does not $\mathfrak{p}$ -fork over A by symmetry of $\mathfrak{p}$ -forking. From this last result, using transitivity again, we get that $B \downarrow_A^{\mathfrak{p}} b$ and $B \downarrow_{Ab}^{\mathfrak{p}} a$. By symmetry $b \downarrow_A^{\mathfrak{p}} B$ and $tp(a/Bb) := q_B(x, b)$ does not $\mathfrak{p}$ -fork over Ab . But $tp(a/Ab)$ is $q(x, b)$ so by induction hypothesis

$$\mathfrak{p}(tp(a/Bb), \phi, \theta, k) = \mathfrak{p}(q(x, b), \phi, \theta, k) \geq n.$$

Since $b \downarrow_A^{\mathfrak{p}} B$, we know we can find some $c' \models tp(c/Ab)$ such that $b \downarrow_{Ac'}^{\mathfrak{p}} B$, $\models \theta(b, c')$ and $\{\phi(x, y)\}_{y \models \theta(y, c')}$ is k -inconsistent. These are the only things

we need c for so we will assume $b \downarrow_{Ac}^b B$. By hypothesis $tp(b/Ac)$ is non-algebraic so by lemma 2.1.2 neither is $tp(b/Bc)$.

Now, for any $b' \models tp(b/Bc)$ we have $\text{p}(q_B(x, b'), \phi, \theta, k) \geq n$, $\phi(x, b') \in q_B(x, b')$ and $\{\phi(x, b')\}_{b' \models \theta(y, c)}$ is k -inconsistent. Therefore,

$$\text{p}(q_B(x, b'), \phi, \theta, k) \geq n$$

which by monotonicity of the p -ranks implies

$$\text{p}(p(x) \cup \{\phi(x, b')\}, \phi, \theta, k) \geq n.$$

By monotonicity and definition of the p -ranks,

$$\text{p}(p, \phi, \theta, k) \geq n + 1$$

□

4.1.1. p -forking and p -Morley sequences. Throughout the development of stability and simplicity theory Morley sequences have been used for characterizing forking in both stability and simplicity theory. In rosy theories we can partially recover such characterization ⁵.

Remark 4.1.4. *Once we have transitivity and symmetry, any p -Morley sequence $\langle a_i \rangle_{i \in I}$ has the property that for any $i \in I$, $tp(a_i/A \cup \{a_j\}_{j \in I, j \neq i})$ does not p -fork over A .*

Proof. The proof given for Morley sequences in simple theories is a straightforward application of symmetry, transitivity and local character. □

Theorem 4.1.5. *If there is a p -Morley sequence over A with $a = a_0$ such that $\bigwedge_i \delta(x, a_i)$ is consistent, then $\delta(x, a)$ does not p -fork over A .*

Proof. Let $\langle a_i \rangle_{i \in \mathbb{N}}$ be a p -Morley sequence over A with $a_0 = a$ and let $\bigwedge_i \delta(x, a_i)$ be consistent, realized by an element so let b . Suppose that $tp(b/Aa_0)$ p -forks over A . By the previous theorem we know that this is witnessed by

$$\text{p}(tp(b/Aa_0), \psi, \theta, k) < \text{p}(tp(b/A), \psi, \theta, k)$$

where ψ and θ are formulas which only depend on the formulas that witness the p -forking of $\delta(x, a)$ ⁶ (see the proof of theorem 3.2.3).

Using an automorphism we get that $tp(b/Aa_n)$ p -forks over A . However,

$$a_{n+1} \downarrow_A^b a_0, \dots, a_n$$

so by corollary 4.1.2 we know that $\delta(x, a_{n+1})$ actually p -forks over $Aa_0a_1 \dots a_n$ and it implies a similar disjunction of formulas which p -divide. By the proof

⁵Partially in the sense that unlike the simple case, it is not true that if we take any p -Morley sequence the conjunction of the resulting formulas would necessarily be consistent –o-minimal theories can witness the failure of this remark–

⁶By this we mean that ψ and θ depend only on the formulas which $\delta(x, a)$ implies and p -divide over A and on the formulas needed to prove the k -inconsistency of such formulas.

of theorem 3.2.3, not only do we know that $tp(b/Aa_0a_1 \dots a_{n+1})$ is a $\mathfrak{p}$ -forking extension of $tp(b/Aa_0a_1 \dots a_n)$ but we also know that for all n ,

$$\mathfrak{p}(tp(b/Aa_0a_1 \dots a_{n+1}), \psi, \theta, k) < \mathfrak{p}(tp(b/Aa_0a_1 \dots a_n), \psi, \theta, k)$$

which is impossible since $\mathfrak{p}(tp(b/Aa_0a_1 \dots a_{n+1} \dots), \psi, \theta, k)$ is finite. $\square$

Theorem 4.1.6. *Suppose $\delta(x, a)$ does not $\mathfrak{p}$ -fork over A . Then there is a $\mathfrak{p}$ -Morley sequence I over A , $a_0 \in I$ such that $\bigwedge_{a_i \in I} \delta(x, a_i)$ is consistent.*

Proof. By extension, if $\delta(x, a)$ does not $\mathfrak{p}$ -fork over A , we can find some b realizing $\delta(x, a)$ such that $tp(b/Aa)$ does not $\mathfrak{p}$ -fork over A . By symmetry, $tp(a/Ab)$ does not $\mathfrak{p}$ -fork over A . We can then construct a $\mathfrak{p}$ -Morley sequence $I = \langle a_i \rangle$ over A where $a_0 = a$ and for all i , $a_i \models tp(a/Ab)$. Such b witnesses the consistency of

$$\bigwedge_{a_i \in I} \delta(x, a_i).$$

$\square$

4.1.2. *Superrosy theories and The U^b -rank.* As with simple theories in some cases we can define a global rank, the U^b -rank, which will share many of the properties that the U -rank has. It will also help us analyze (in chapter 3) the relation between $\mathfrak{p}$ -forking and usual forking in supersimple theories.

Definition 4.1. *We define the U^b -rank inductively as follows. Let $p(x)$ be a type over some set A . Then,*

- (1) $U^b(p(x)) \geq 0$ if $p(x)$ is consistent.
- (2) For any ordinal α , $U^b(p(x)) \geq \alpha + 1$ if there is some tuple a and some type $q(x, a)$ over Aa such that $q(x, a) \supset p(x)$, $U^b(q(x, a)) \geq \alpha$ and $q(x, a)$ $\mathfrak{p}$ -forks over A .
- (3) For any λ limit ordinal, $p(x) \geq \lambda$ if and only if $p(x) \geq \sigma$ for all $\sigma < \lambda$.

Remark 4.1.7. *Using extension to get from $\mathfrak{p}$ -forking to $\mathfrak{p}$ -dividing, one can easily verify that we can replace condition (2) by the following statement:*

- (2') For any ordinal α , $U^b(p(x)) \geq \alpha + 1$ if there is some $p(x, a) \supset p(x)$ and some c such that
 - $U^b(p(x, a)) \geq \alpha$.
 - $tp(a/Ac)$ is non-algebraic.
 - $\{p(x, a')\}_{a' \models tp(a/Ac)}$ is k -inconsistent for some k .

Definition 4.2. *A theory T is superrosy if given any set A in a model of T and a type $p(x) \in S(A)$, $U^b(p(x))$ is defined.*

Equivalently, a theory is superrosy if there are no infinite forking chains of types.

In the case of supersimple theories (theories where the SU -rank of any type is defined), one of the most used properties is that this rank satisfies Lascar's inequalities. With the properties we have proved for $\mathfrak{p}$ -forking, the

proof of the Lascar's inequalities for simple theories (see [Wag00]) works in our context.

Theorem 4.1.8. Lascar's Inequalities: *Whenever the U^b -rank is defined, it satisfies the following inequality:*

$$U^b(tp(a/bA)) + U^b(tp(b/A)) \leq U^b(tp(ab/A)) \leq U^b(tp(a/bA)) \oplus U^b(tp(b/A))$$

Proof. See [Wag00] theorem 5.1.6. $\square$

Finally, we want to give an alternative definition of U^b -rank which is closer to the work in hyperimaginaries and because of this it will prove very useful when comparing our U^b and the usual SU-rank in supersimple theories.

Definition 4.3. *We define the U_*^b -rank on types inductively as follows. Let $p(x)$ be a type over some set A . Then,*

- (1) $U_*^b(p(x)) \geq 0$ if $p(x)$ is consistent.
- (2) For any ordinal α , $U_*^b(p(x)) \geq \alpha + 1$ if there is some $p(x, a)$ and some set $B \supset A$ such that
 - $p(x, a) \supset p(x)$.
 - $U_*^b(p(x, a')) \geq \alpha$ for any $a' \models tp(a/B)$ and $tp(a/B)$ non-algebraic.
 - There is a $k \in \mathbb{N}$ such that for any distinct $a_1, a_2, \dots, a_k \models tp(a/B)$, $\bigcup_{i=1}^k p(x, a_i)$ is a b -forking extension of $p(x, a_1)$ ⁷.
- (3) For any λ limit ordinal, $p(x) \geq \lambda$ if $p(x) \geq \sigma$ for all $\sigma < \lambda$

The main part of proving the two definitions are equivalent is the following lemma, which in itself is a weak amalgamation for b -independent extensions of a given type. It is far away, however, from the Independence Theorem that characterizes simple theories. As mentioned in the introduction, in chapter 4 we will talk about which of these amalgamation theorems can be true in a general rosy theory.

Lemma 4.1.9. *Let A, B be supersets of some set C such that $A \downarrow_C^b B$. Let p_A and p_B be non- b -forking extensions to A and B of the same type p over C . Then there is some $B' \models tp(B/C)$, $B' \downarrow_C^b A$ such that $p_{B'} \cup p_A$ does not b -fork over C .*

Proof. Let a, b be two elements such that $a \models p_A$ and $b \models p_B$. By hypothesis, $a \downarrow_C^b A$ and $b \downarrow_C^b B$. Since a and b satisfy the same type over C , there is an automorphism σ fixing C such that $\sigma(b) = a$; let $B'' := \sigma(B)$ so $a \downarrow_C^b B''$. Using symmetry and extension, we can find some $B' \models tp(B''/Ca)$ such that $B' \downarrow_C^b Aa$ (so in particular $a \models p_{B'}$ and $B' \downarrow_C^b A$). Using transitivity and symmetry once more we get $a \downarrow_A^b CB'$ and by transitivity $a \downarrow_C^b AB'$.

By construction, $B' \models tp(B/C)$ and $B' \downarrow_C^b A$. Finally, $a \models p_A \cup p_{B'}$ and $a \downarrow_C^b AB'$. $\square$

⁷If $\bigcup_{i=1}^k p(x, a_i)$ is inconsistent it is considered to be a b -forking extension of $p(x, a_1)$

Claim 4.1.10. *For any complete type p ,*

$$U^b(p) = U_*^b(p).$$

Proof. We will prove that for any complete type p , $U^b(p) \geq U_*^b(p)$ and $U^b(p) \leq U_*^b(p)$.

To prove that $U^b(p) \geq U_*^b(p)$ we will prove that for any ordinal α , $U^b(p) \geq \alpha$ implies $U_*^b(p) \geq \alpha$. For $\alpha = 0$ and or when α is a limit ordinal, the induction follows from the definitions; the successor case follows from remark 4.1.7.

As for the other direction, we need to prove that

$$(1) \quad \text{if } U_*^b(p) \geq \alpha \text{ then } U^b(p) \geq \alpha.$$

We will again do an induction on α . For $\alpha = 0$ or α a limit ordinal the induction is immediate. For the successor case, let us assume that (2.1) is true for all $\alpha \leq \sigma$ and that $U_*^b(p) \geq \sigma + 1$. By definition we can find some a and some $B \supset A$ such that $p(x, a)$ is an extension of $p(x)$, $U_*^b(p(x, a)) \geq \sigma$ and there is some k such that for any $a_1, a_2, \dots, a_k \models tp(a/B)$, $U_*^b(\cup_{i=1}^k p(x, a_i)) < U_*^b(p(x, a))$. If $p(x, a)$ $\mathfrak{p}$ -forks over A we get the conclusion of the claim by definition of U^b -rank (after extending the types so we get $\mathfrak{p}$ -division without changing the ranks). We will show that this is the only possible case.

Suppose that $p(x, a)$ does not $\mathfrak{p}$ -fork over A . Let $p(x, a, B)$ be a non- $\mathfrak{p}$ -forking extension of $p(x, a)$ to Ba . Since $\mathfrak{p}$ -forking is preserved by automorphisms, we know that for any $a' \models tp(a/B)$ $p(x, a', B)$ is a non- $\mathfrak{p}$ -forking extension of $p(x, a')$ and

$$p(x, a', B) \upharpoonright B = p(x, a, B) \upharpoonright B := p(x, B).$$

Given any such a' , $p(x, a', B)$ does not $\mathfrak{p}$ -fork over A by transitivity and by monotonicity $p(x, a', B)$ is a non- $\mathfrak{p}$ -forking extension of $p(x, B)$.

Now, applying lemma 4.1.9 k times, we can find a sequence $\langle a_1, a_2, \dots, a_k \rangle$ with $a = a_1$ such that for any $i \leq k$, $a_i \models tp(a/B)$ and for all $i, j, i \neq j$, $a_i \downarrow_B^{\mathfrak{p}} a_j$ and $\cup_{i=1}^k p(x, a_i, B)$ is a non- $\mathfrak{p}$ -forking extension of $p(x, B)$. But restricting to $A\bar{a}_i$ we get that $\cup_{i=1}^k p(x, a_i)$ is a non- $\mathfrak{p}$ -forking extension of $p(x)$ and therefore a non- $\mathfrak{p}$ -forking extension of $p(x, a_1)$, a contradiction. $\square$

5. $\mathfrak{p}$ -FORKING IN SIMPLE AND O-MINIMAL THEORIES

In this chapter we will look at theories for which independence notions have been studied by other authors. In particular, we are interested in analyzing the behavior of $\mathfrak{p}$ -forking inside simple and o-minimal theories which are, as we mentioned in the introduction, the two classes of theories where the independence notion has been a main tool for the development of the respective theories.

5.1. Simple and Stable cases. Given any theory, it is clear by the definition of $\mathfrak{p}$ -forking that any formula which $\mathfrak{p}$ -divides over some set A $\mathfrak{p}$ -forks over A ; Also by definition, for any type $p(x)$, any two sets of formulae Δ , Π and any $k \in \mathbb{N}$, $\mathfrak{p}(p, \Delta, \Pi, k) \leq D(p, \Delta, k)$. So all simple theories are rosy theories and $\mathfrak{p}$ -forking is a stronger notion than forking in the sense that for any sets A, B, C , $A \perp_C B \Rightarrow A \perp_C^{\mathfrak{p}} B$. As for the converse, we will prove in this section that in all known examples of simple theories the two notions agree. In fact we prove that the two notions being equal is implied by the (strong) stable forking conjecture.

5.1.1. $\mathfrak{p}$ -forking inside stable theories. We will prove that in a stable theory $\mathfrak{p}$ -forking and forking are the same. The following theorem will be the key to our proof.

Theorem 5.1.1. *Inside a stable theory T , if a type p forks over A , then there is a non-forking extension q of p , a formula δ and a non-algebraic type π (in $\mathcal{L}^{eq}$) such that $\{\delta(x, a)\}_{a \models \pi}$ is k -inconsistent and $\delta(x, a) \in q$ for some $a \models \pi$.*

Proof. Let $\phi(x, b_0)$ be a formula implied by p which divides over A , and let $\{\phi(x, b_i)\}$ be the set of k -inconsistent formulas which witness the dividing, with $\langle b_i \rangle$ an A indiscernible sequence. Since we are working inside a stable theory, we can assume $k = 2$. Let $\Phi(x, y) = \{\phi(x, y), x = y\}$, $\pi_1 = tp(b_0/A)$.

Since we are only looking for a non-forking extension of p , we can assume p to be a complete type in $S(\text{acl}(A))$ so that $(R, d)_{\Phi}(p) = (\alpha, 1)$. Let ϕ_0 be a Φ formula in p with the same Φ -rank and degree as p . Then $\phi_0 \wedge \phi(x, b_0)$ we be an A -dividing Φ -formula such that $(R, d)_{\Phi}(p) = (R, d)_{\Phi}(\phi_1)$; we can assume that $(R, d)_{\Phi}(p) = (R, d)_{\Phi}(\phi(x, b_0))$.

To complete the proof we will need some sort of pre- $\mathfrak{p}$ -dividing in a definable class of formulas in p (or in a non-forking extension). We suspect this may have been done in the theory of canonical bases in stability theory. However, we were not able to find a statement in the literature with exactly the properties we needed so we will prove the following claim.

Claim 5.1.2. *Let p be a type in $S(\text{Ab}_0)$ such that $(R, d)_{\Phi}(p) = (\alpha, 1)$. Let $\phi(x, b_0) \in p$ such that $(R, d)_{\Phi}(\phi(x, b_0)) = (R, d)_{\Phi}(p)$. Then we can find formula $\eta(x, [b])$ and a type q such that*

- (1) $\eta(x, [b])$ belongs to $p(x, b_0)$,
- (2) $R_{\Phi}(\eta(x, [b])) = \alpha$,
- (3) For any $[a], [b]$ such that $[a], [b] \models q$ and $[a] \neq [b]$,

$$R_{\Phi}(p(x) \cup (\eta(x, [b]) \wedge \eta(x, [a]))) < R_{\Phi}(p).$$

If, besides, there is an infinite sequence $\langle b_i \rangle$ of elements satisfying $tp(b_0/A)$ such that $\{\phi(x, b_i)\}$ is 2-inconsistent, then $tp([b]/A)$ is non-algebraic.

Proof. As in (see [Zie98] 2.2), given two definable classes $\mathbb{F}$ and $\mathbb{G}$, we define

$$(2) \quad \mathbb{F} \subset_{\Phi, \alpha} \mathbb{G} \text{ if } R_{\Phi}(\mathbb{F} \setminus \mathbb{G}) < \alpha$$

and

$$(3) \quad \mathbb{F} \cong_{\Phi, \alpha} \mathbb{G} \text{ if } R_{\Phi}(\mathbb{F} \Delta \mathbb{G}) < \alpha.$$

Given formulas $\psi_1(x), \psi_2(x)$, we will abbreviate $\psi_1(\mathbb{C}) \subset_{\Phi, \alpha} \psi_2(\mathbb{C})$ by $\psi_1(x) \subset_{\Phi, \alpha} \psi_2(x)$.

Let ϕ' be the formula $\phi(x, y)$ with variable y and parameters in x . Let $\pi(y)$ be $tp(b_0/A)$. For any $b \models \pi$, let $\psi(b, y)$ be the ϕ definition of $p \cup \{\phi(x, b)\}$ (which is a ϕ' -formula over Ab). Then $\models \psi(b, c)$ if and only if $\phi(x, b) \subset_{\Phi, \alpha} \phi(x, c)$, for any c . By using an automorphism between b and any other $a \models \pi$, we know that $\psi_a(y) = \psi(a, y)$. Let

$$E_1(z, y) := \psi(z, y)$$

For any $a \models \pi$, the Φ -degree of $\phi(x, a)$ is 1 so

$$\begin{aligned} \psi(a, b) &\iff \phi(x, b) \subset_{\Phi, \alpha} \phi(x, a) \\ &\iff R_{\Phi}(\phi(x, b) \wedge \neg \phi(x, a)) < \alpha \\ &\iff R_{\Phi}(\phi(x, b) \wedge \phi(x, a)) = \alpha \\ &\iff R_{\Phi}(\neg \phi(x, b) \wedge \phi(x, a)) < \alpha \\ &\iff \psi(b, a) \\ &\iff \phi(x, a) \cong_{\Phi, \alpha} \phi(x, b) \end{aligned}$$

So $\pi(z) \wedge \pi(y) \vdash "E_1 \text{ is an equivalence relation}"$ and there is some finite $\pi_1 \subset \pi$ such that $\pi_1(z) \wedge \pi_1(y) \wedge E_1(z, y)$ is an equivalence relation E_2 . However, we may have more points than we want in each equivalence class so we restrict it a bit more by finding a finite $\pi_2 \subset \pi$ such that for some (and because π is a complete type over A for all) $b \models \pi$,

$$(R, d)_{\phi'}(\pi_2(y) \cup \{E_2(b, y)\}) = (R, d)_{\phi'}(\pi(y) \cup \{E_2(b, y)\}).$$

Let

$$E(z, y) := \pi_2(z) \wedge \pi_2(y) \wedge E_2(z, y).$$

$E(z, y)$ is contained in $\pi(z) \wedge \pi(y) \wedge E(z, y)$ so for any $a, b \models \pi$

$$(4) \quad E(a, b) \iff E_1(a, b) \iff R_{\Phi}(\neg \phi(x, a) \wedge \phi(x, b)) < \alpha$$

and, by additivity of R_{Φ} -rank in stable theories,

$$(5) \quad \neg E(a, b) \Rightarrow \neg E_1(a, b) \Rightarrow R_{\Phi}(\phi(x, a) \wedge \phi(x, b)) < \alpha;$$

let $[b]$ be the E -equivalence class of b .

We will now work with generic points to make calculations easier. Let b satisfy a non-forking extension of $tp(b_0/A[b_0])$ such that $b \perp_{A[b_0]} b_0$. Let $c \models p(x, b_0)$ such that $b \perp_{Ab_0}^b c$.

Let $\eta(x)$ be the ϕ' definition of $tp(b/Ab_0)$ (and because $tp(b/Ab_0)$ is a non-forking extension of $tp(b/A[b_0])$ all the parameters of η are in $acl(A[b_0])$)

so $\eta = \eta(x, [b_0])$ is over $\text{acl}(A[b_0])$. In particular, we know that $b \models tp(b_0/A[b_0])$, $c \models p(x, b_0)$ and $\psi(b_0, y) \in tp(b/Ab_0)$. It follows that $\eta(x, [b_0]) \in p$ (see [Pil96] lemma 2.8).

We can now prove that such $\eta(x, [b_0])$ satisfies condition 3. Suppose not and let $[b] \neq [a]$ such that $R_\Phi(p(x) \cup (\eta(x, [b]) \wedge \eta(x, [a]))) = \alpha$. We extend $p(x)$ to a non-forking type $p'(x)$ over a model M .

Let a', b' be elements in $[a]$ and $[b]$ respectively such that $a' \downarrow_{[a]} A$ and $b' \downarrow_{[b]} A$. Let $c \models p(x) \cup (\eta(x, [b]) \wedge \eta(x, [a]))$ such that $c \downarrow_M a'b'$; such c satisfies the ϕ' definition of $tp(a'/A[a])$ and $tp(b'/A[b])$. By [Pil96] 2.8 again and non-forking, both $b', a' \models \phi(c, y)$. But $[b'] = [b] \neq [a] = [a']$ so $b' \not\equiv_{\Phi, \alpha} a'$. By definition,

$$(R, d)_\Phi(\phi(x, b') \wedge \phi(x, a')) < (\alpha, 1)$$

which contradicts $c \downarrow_M a'b'$.

Finally, if $\langle b_i \rangle$ is an infinite sequence of elements satisfying π such that $\{\phi(x, b_i)\}$ is 2-inconsistent, then each b_i in the sequence belongs to a different class $[b_i]$ which proves the last statement of the claim. $\square$

We can prove the theorem using lemma 4.1.9 and a careful induction on the ranks. However, this proof may not work in the general context of simple theories with stable forking conjecture and it does not provide a specific $\mathfrak{p}$ -forking formula which is always nice to have. We give a different proof which, even though it is more technical, provides an algorithm to construct the $\mathfrak{p}$ -dividing formula and it is a good insight of what happens, for example, in algebraically closed fields.

We know that $\eta(x, [b_0])$ in the claim forks over A so there must be some formula in the η -type of $p(x, b_0) \upharpoonright \text{acl}(A[b_0])$ which divides over A . We can take this new formula instead of $\phi(x, b_0)$ in the statement of the theorem and assume that $\phi(x, b)$ itself satisfies the conclusion of the claim or that $\eta(x, [b_0])$ is a Φ -formula.

Given any two $\beta, d \in N$, the statement “There is an infinite subfamily I of $b_i \models \pi$ such that $(R, d)_\Phi \{\phi(x, b_i)\}_{b_i \in I} \geq (\beta, d)$ ” is type definable. If $\{\phi(x, b)\}_{b \models tp(b_0/A)}$ is not k -inconsistent there is an infinite subfamily which is consistent. Let (β_1, d_1) be the maximal Φ -rank and -degree that an infinite subfamily of $\{\phi(x, b)\}_{b \models tp(b_0/A)}$ can achieve⁸. Let B be an infinite set of elements in $tp(b_0/A)$ such that

$$R_\Phi(\{\phi(x, b')\}_{b' \in B}) = (\beta_1, d_1).$$

By using an automorphism we can assume that this set includes $\phi(x, b_0)$. Note that by the previous claim β_1 must be less than α .

⁸So that for any infinite subfamily I' of $\{b \mid b \models tp(b_0/A)\}$,

$$(R, d)_\Phi\left(\bigcup_{b \in I'} \{\phi(x, b)\}\right) \leq (\beta_1, d_1).$$

Such a maximum exists by compactness

The rest of the proof will consist of narrowing down the set of sentences we will consider, and maybe working with non-forking extensions.

We will define a sequence of formulas $\{\theta_i(x)\}$ inductively. Let $\phi_1(x, b) = \phi(x, b)$, q_1 be $\{\phi(x, b)\}_{b \in B}$ so $(R, d)_\Phi(q_1) = (\beta_1, d_1)$.

By the local property of the rank, there is a finite conjunction of $\phi(x, b_i) \in q_1$ (which we will call θ_1) such that $(R, d)_\Phi(\theta_1) = (R, d)_\Phi(q_1)$; so for any b_i , $\phi_1(x, b_i) \wedge \theta_1$ is a forking extension of $\phi_1(x, b_i)$. By extension, $p(x, b_i) \cup \{\neg\theta_1\}$ is a non-forking extension $p_2(x, b_i^2)$ of $p(x, b_i)$. Let $r_1(x)$ be a non-forking extension of θ_1 to a model M and let $\gamma(y)$ be its ϕ -definition. By compactness we can assume B is very big (big enough to use Erdos-Rado), and by Erdos-Rado we can also assume it contains some infinite M -indiscernible subset $< b_j >_{j \in J}$ (as usual we can get $b_{j_0} = b_0$). For all $j \in J$ let $\phi_2(x, b_j^2) = \phi(x, b_j) \wedge \neg\theta_1(x)$ which is consistent (and in fact a non-forking extension of $\phi(x, b_j)$) for all $j \in J$.

Recall that $q_1(x)$ had the same Φ rank and degree as θ_1 so that $b_{j_0} = b_0 \models \gamma(y)$. We will say that $b_i^2 \models \gamma(y)$ whenever the projection $b_j \models \gamma(y)$.

Let $\pi_2 = tp(b_0^2/M) \cup \gamma(y)$. This type is satisfied by all the b_i^2 's and is therefore non algebraic.

For any infinite subfamily $s(x)$ of $\{\phi_2(x, b_i^2)\}_{b_i^2 \models \pi_2}$, we know that $s(x)$ is of the form $\{\phi(x, b_i)\}_{i \in J} \cup \{\neg\theta_1(x)\}$ where J is some index set such that $b_i \models tp(b_0/M)$ for any $i \in J$. By maximality of $(R, d)_\Phi(q_1)$ we have $(R, d)_\Phi(r) \leq (\beta_1, d_1)$. If $\{\phi_2(x, b_i^2)\}_{b_i^2 \models \pi_2}$ is not k -inconsistent for any k , we can once more find some consistent subset $I_2 \subset \pi_2(\mathcal{C})$ such that $(\bigwedge_{i \in I_2} \phi_2(x, b_i^2))$ has maximum $(R, d)_\Phi$ in the lexicographical order; say $(R, d)(\bigwedge_{i \in I_2} \phi_2(x, b_i^2)) = (\beta_2, d_2) \leq (\beta_1, d_1)$.

Recall that for $i \in I_2$, $b_i^2 \models \gamma(y)$ (the ϕ definition of θ_1) so

$$(R, d)_\Phi(\{\phi(x, b_i)\}_{i \in I_2} \cup \theta_1(x)) = (\beta_1, d_3)$$

for some d_3 . If $\beta_1 = \beta_2$ we get, using the fact that θ_1 is a Φ -formula, that

$$\begin{aligned} \deg_\Phi(\{\phi(x, b_i)\}_{i \in I_2}) &= \deg_\Phi(\{\phi_2(x, b_i^2)\}_{i \in I_2}) + \deg_\Phi(\{\phi(x, b_i)\}_{i \in I_2} \cup \theta_1(x)) \\ &\Rightarrow d_1 \geq d_2 + d_3 \\ &\Rightarrow d_2 < d_1. \end{aligned}$$

So $(\beta_2, d_2) <^{lex} (\beta_1, d_1)$. Let $q_2 = \{\phi(x, b_i^2)\}_{i \in I_2}$ and θ_2 be a conjunction of $\phi_2(x, b_i^2)$ with the same Φ -rank, degree as q_2 .

We can repeat the above construction as long as

$$\{\phi_n(x, b_i^n)\}_{b_i^n \models \pi_n}$$

is not k -inconsistent for any k ; by doing this we would get a sequence $\langle (R, d)_\Phi(\theta_i) \rangle_{i \leq n}$ which is strictly decreasing in the lexicographical order.

Since both the Φ rank and degrees are ordinals (and therefore well ordered) at some point we must get some N such that $\{\phi_N(x, b_i^N)\}_{b_i^N \models \pi_N}$ is k -inconsistent and by construction, such $\phi_N(x, b_0^N)$ would belong to a non-forking extension of $p(x, b_0)$. $\square$

Corollary (to the proof). *If in some model of a simple theory, some type p has a stable formula which forks over A , then there is a non-forking extension q of p , a stable formula δ and a type π such that $\{\delta(x, a)\}_{a \models \pi}$ is k -inconsistent and $\delta(x, a) \in q$ for some $a \models \pi$*

Proof. If ϕ is stable, then so is the formula η defined in the claim (it is equivalent to a conjunction of δ formulas) and so are all the formulas defined in the rest of the proof. $\square$

Definition 5.1. *A theory T has the strong stable forking property if whenever p is a type over some B subset of $\mathcal{C}$ and p forks over a set A (not necessarily contained in B), then this is witnessed by some instance of a stable formula in p . We will say that T satisfies the weak stable forking property if we require B to be a model of T and A to be a subset of B .*

Corollary 5.1.3. *In any stable theory T , if $M \models T$ is a model of T , A is a subset of M , $q \in S(M)$ is a complete type and $p = q \upharpoonright A$, then q is a non-forking extension of p if and only if it is a non- $\mathfrak{p}$ -forking extension. In fact, this conclusion is true for any simple theory T for which the strong stable forking property holds⁹ (which includes all known examples of simple theories).*

Proof. The left to right implication is immediate since $\mathfrak{p}$ -forking is a stronger notion than forking. For the other side, if q forked over A there would be some $r(x, b')$ a non-forking extension of q , a formula δ and a non-algebraic type π such that $\{\delta(x, b)\}_{b_i \models \pi}$ is k -inconsistent, $b' \models \pi$, $\delta(x, b') \in r(x, b')$. Let $\{b_i\}_{i \in N}$ be an infinite set of A -conjugates of b' . By monotonicity,

$$\mathfrak{p}(p(x) \wedge \delta(x, b_i), \delta, \pi, k) \geq \mathfrak{p}(r(x, b_i), \delta, \pi, k).$$

Conjugating over A we get $\mathfrak{p}(r(x, b'), \delta, \pi, k) = \mathfrak{p}(r(x, b_i), \delta, \pi, k)$. By definition,

$$\mathfrak{p}(p, \delta, \pi, k) \geq \mathfrak{p}(r(x, b'), \delta, \pi, k) + 1.$$

But any non-forking extension is a non- $\mathfrak{p}$ -forking extension so

$$\mathfrak{p}(r(x, b'), \delta, \pi, k) = \mathfrak{p}(q, \delta, \pi, k)$$

so

$$\mathfrak{p}(p, \delta, \pi, k) < \mathfrak{p}(q, \delta, \pi, k).$$

$\square$

⁹Even though the proof is done assuming the strong stable forking property, we think the weak stable forking conjecture might be enough.

5.1.2. Supersimple theories. In this section we shall prove that for supersimple theories the forking relation is equivalent to the $\mathfrak{p}$ -forking one. It is clear that for any type p , $U(p) \geq U^{\mathfrak{p}}(p)$; so in supersimple theories both the U and the $U^{\mathfrak{p}}$ are defined and they both characterize forking and $\mathfrak{p}$ -forking respectively. Therefore, it is enough to prove the inequality assuming both these ranks are ordinal valued.

Theorem 5.1.4. *For any type p*

$$U(p) = U^{\mathfrak{p}}(p)$$

Proof. The U -rank is always bigger than the $U^{\mathfrak{p}}$ -rank and we proved the equivalence between $U_*^{\mathfrak{p}}$ and $U^{\mathfrak{p}}$, so it is enough to show that for any ordinal α and any type p ,

$$U(p) = \alpha \Rightarrow U_*^{\mathfrak{p}}(p) \geq \alpha.$$

We will do an induction on $U(p)$. For $\alpha = 0$ or a limit ordinal it is clear. Let us assume then that $U(p) = \alpha = \sigma + 1$ and that for any $\lambda \leq \sigma$ and any type p , $U(p) = \lambda \Rightarrow U_*^{\mathfrak{p}}(p) \geq \lambda$. But $U(p) \geq U_*^{\mathfrak{p}}(p)$ so our induction hypothesis is equivalent to $U(p) = \lambda \Rightarrow U_*^{\mathfrak{p}}(p) = \lambda$.

Let p be a type over some set A such that $U(p) = \sigma + 1$. By definition, we can extend p to some type $q(x, C)$ over $C \supset A$ such that $U(q) = \sigma$ and q forks over A . Let a' be the canonical base of q over A ¹⁰ and let $p(x, a') = q \upharpoonright Aa'$. Now, by elimination of hyperimaginaries in supersimple theories (see [BPW01]), we can assume a' is a tuple in the model; by definition of the canonical base we know that for any two $a_1, a_2 \models tp(a'/A)$, $p(x, a_1) \cup p(x, a_2)$ is a forking extension of $p(x, a_1)$ and by [Kim96] theorem 3.3.6 we have that $q(x)$ does not fork over Aa' . Now, in supersimple theories this translates into the following statement. For any $a_1, a_2 \models tp(a'/A)$,

$$\sigma = U(q) = U(p(x, a_1)) > U(p(x, a_1) \cup p(x, a_2)).$$

By induction hypothesis,

$$\sigma = U_*^{\mathfrak{p}}(p(x, a_1)) > U_*^{\mathfrak{p}}(p(x, a_1) \cup p(x, a_2)).$$

By definition of the $U_*^{\mathfrak{p}}$ we get $U_*^{\mathfrak{p}}(p) \geq \sigma + 1 = U(p) \geq U^{\mathfrak{p}}(p) = U_*^{\mathfrak{p}}(p)$. So $U_*^{\mathfrak{p}}(p) = \sigma + 1 = U(p)$. $\square$

5.2. O-minimal: We shall prove that in o-minimal theories the global $U^{\mathfrak{p}}$ rank is defined and equal to the usual dimension. This will prove that $\bigcup^{\mathfrak{p}}$ is precisely the usual independence relation. Let M be the model of some o-minimal theory.

Claim 5.2.1. *For any definable $A \subset M^k$, if $\dim(A) = n$ then $U^{\mathfrak{p}}(A) \leq n$*

¹⁰The construction of such hyperimaginary would be made by defining an equivalence relation just for extensions of $p(x)$ (looking at A as a fixed set of parameters). We might have to extend $p(x)$ to a type over a model containing A , but all this can be done by the extension and transitivity properties.

Proof. Suppose otherwise and let i be the smallest integer such that there is a definable A , $\dim(A) = i$ and $\text{U}^b(A) \geq i+1$, let A be definable over some finite set B by some formula ϕ , and let $A \subseteq M^k$. We can then find some “generic” point (in the sense of the U^b -rank) $a \in A$ such that $\text{U}^b(tp(a/B)) \geq i+1$. By minimality of i , $\dim(tp(a/B)) \geq i$ (otherwise we could find some definable subset contained in A for which a is dimension-generic). By the theory of o-minimal sets (see [vdD98]) there is some open subset A' with $a \in A' \subset A$ such that we can find a projection from M^k to M^i which is a definable homomorphism from A' into some open subset of M^i . But we know that $\text{U}^b(A') \geq i+1$ and $\dim(A') = i$, both of which are preserved by definable homomorphisms so we may assume that $A \subseteq M^i$.

By definition of the U^b -rank, there is some formula $\delta(x, y)$, an element b_0 and some $C \supset B$ such that

- $tp(b_0/C)$ is non algebraic
- $\text{U}^b(tp(a/B) \cup \delta(x, b_0)) \geq i$ and
- $\{\delta(x, b')\}_{b' \models tp(b_0/C)}$ is k -inconsistent.

We will first do the case where $i = 1$. Let $\langle b_i \rangle_{i \in \mathbb{N}}$ be some indiscernible sequence in $tp(b_0/C)$ and A_j be the set defined by $\delta(x, b_j)$. By hypothesis $\text{U}^b(A_j) = 1$ and $A_j \subset M^1$ so all of the A_j 's are infinite subsets of an o-minimal model. By definition of o-minimality, each of the A_j 's must contain an interval. We could therefore (uniformly) define A'_j as the first interval contained in A_j and we can define the subset $X \subset M$ consisting of left endpoints of the A'_j . By k -inconsistency, this set cannot contain an interval and it must be infinite (not more than $k-1$ of the A_j 's can have the same left endpoint) contradicting o-minimality.

For a general i , note that by minimality of i we have that $\dim(\phi(x) \wedge \delta(x, b')) \geq i$ for any $b' \models tp(b/C)$. But it is a definable subset of M^i so the projection to each of the coordinates must be infinite and definable. Taking the projection to the first coordinate

$$\phi_0(x_0, b') := \exists x \phi(x) \wedge \delta(x, b') \wedge \pi_0(x) = x_0$$

we have a definable set in M^1 so by the $i = 1$ case the set of formulas $\{\phi_0(x_0, b')\}_{b' \models tp(b_0/C)}$ cannot be k -inconsistent. By definition of k -inconsistency we can find an infinite subsequence $\langle b_j \rangle_{j \in J}$ of $\{b'\}_{b' \models tp(b_0/B)}$ such that

$$\bigwedge_{b_i} \phi_0(x_0, b_i)$$

is consistent; say it is realized by a_0 . Let $q_0(y)$ be the (non algebraic) type $tp(b_0/B) \cup \{\phi_0(a_0, y)\}$ and consider $\{\delta(x, b')\}_{b' \models q_0}$. We repeat the procedure with each of the projections (we always get non algebraic types) and inductively define a point in M^i $a := (a_0, \dots, a_{i-1})$ which realizes infinitely many of the $\delta(x, b')$'s, contradicting k -inconsistency. $\square$

The other inequality is much easier. If $A \subset M^k$ and $\dim(A) = n$ then there is some open subset A' of A and a definable projection π from M^k to

M^n such that A' gets sent to an open subset U of M^n and $\pi : A' \rightarrow U$ is a homomorphism. Thus, $\dim(U) = n$. This implies that the projection to each of its coordinates is infinite, but $x_i = a$ is clearly a $\mathfrak{p}$ -forking formula for any $i \leq n$ which proves $U^{\mathfrak{p}}(U) \geq n$ so $U^{\mathfrak{p}}(A') \geq n$. By monotonicity of the rank, $U^{\mathfrak{p}}(A) \geq n$.

REFERENCES

- [BPW01] Steven Buechler, Anand Pillay, and Frank Wagner. Supersimple theories. *J. Amer. Math. Soc.*, 14(1):109–124 (electronic), 2001.
- [CW02] Enrique Casanovas and Frank Wagner. The free roots of the complete graph. *Submitted for publication*, 2002.
- [Hod93] Wilfrid Hodges. *Model theory*, volume 42 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1993.
- [Hod97] Wilfrid Hodges. *A shorter model theory*. Cambridge University Press, Cambridge, 1997.
- [Kim96] Byunghan Kim. *Simple First Order Theories*. PhD thesis, University of Notre Dame, 1996.
- [KP97] Byunghan Kim and Anand Pillay. Simple theories. *Ann. Pure Appl. Logic*, 88(2-3):149–164, 1997. Joint AILA-KGS Model Theory Meeting (Florence, 1995).
- [KP01] Byunghan Kim and A. Pillay. Around stable forking. *Fund. Math.*, 170(1-2):107–118, 2001. Dedicated to the memory of Jerzy Łoś.
- [Pil96] Anand Pillay. *Geometric stability theory*, volume 32 of *Oxford Logic Guides*. The Clarendon Press Oxford University Press, New York, 1996. Oxford Science Publications.
- [She90] S. Shelah. *Classification theory and the number of nonisomorphic models*, volume 92 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam, second edition, 1990.
- [vdD98] Lou van den Dries. *Tame topology and o-minimal structures*, volume 248 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1998.
- [Wag00] Frank O. Wagner. *Simple theories*, volume 503 of *Mathematics and its Applications*. Kluwer Academic Publishers, Dordrecht, 2000.
- [Zie98] Martin Ziegler. Introduction to stability theory and Morley rank. In *Model theory and algebraic geometry*, volume 1696 of *Lecture Notes in Math.*, pages 19–44. Springer, Berlin, 1998.